



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

LYCOPODIACEAE

Lycopodium adpressum (Chapm.) Lloyd & Underw.—Southold in a sandy bog.
No. 3455.

L. obscurum L.—Moist woods at Orient and Southold.

(To be continued)

NEW COMBINATIONS FOR PHANEROGAMIC NAMES

By J. C. ARTHUR

In order to secure uniformity in citing the names of hosts for species of Uredinales the following new combinations are proposed. So far as the writer can ascertain these combinations have not been made before, and in coming to this conclusion he has had the kindly assistance of a number of correspondents.

Cnidoscolus urens (L.) comb. nov. (*Jatropha urens* L. Sp. Pl. 1007. 1753). A common plant of tropical America, bearing *Uromyces oaxacanus* Diet. & Holw.

Adenoropium angustifolium (Griseb.) comb. nov. (*Jatropha angustifolia* Griseb.; Goett. Nachr. 171. 1865). A Cuban species bearing the imperfectly known rust *Uredo jatrophiicola* Arth.

Vincetoxicum bifidum (Hemsl.) comb. nov. (*Gonolobus bifidus* Hemsl., Biol. Centr. Am. Bot. 2: 330. 1879).

Vincetoxicum erianthum (Decaisne) comb. nov. (*Gonolobus erianthus* Decaisne; DC. Prodr. 8: 592. 1844).

Vincetoxicum uniflorum (H.B.K.) comb. nov. (*Gonolobus uniflorus* H.B.K. Nov. Gen. Sp. 3: 207. 1818). These three Mexican species of *Vincetoxicum*, belonging to the Asclepiadaceae, bear the very common tropical rust *Puccinia obliqua* Berk. & Curt.

Sphaeralcea arcuata (Greene) comb. nov. (*Malvastrum arcuatum* Robinson; A. Gray, Synop. Fl. N. Am. 1¹: 311. 1878).

Sphaeralcea fasciculata (Nutt.) comb. nov. (*Malva fasciculata* Nutt.; T. & G. Flora N. Am. 1: 225. 1838). These two Californian species belonging to Malvaceae bear the common western rust *Puccinia Sherardiana* Körn.

Madronella viridis (Jepson) comb. nov. (*Monardella viridis* Jepson, Flora W. Mid. Calif. 465. 1901). A plant of western California bearing *Puccinia Monardellae* Dudl. & Thomp., a distinctively Californian rust.

Coleosanthus megalodontus (Greenm.) comb. nov. (*Brickellia megalodonta* Greenm. Proc. Am. Acad. 40: 34. 1904). A Mexican plant bearing the rust *Puccinia Brickelliae* Peck.

PURDUE UNIVERSITY,
LAFAYETTE, INDIANA

SHORTER NOTES

NOTES ON *HEMEROCALLIS*, II.—A previous note (Amer. Mid. Nat. 1914-15) dealt with the nomenclature, specific description, and the distribution of the North American members of this genus, *H. fulva* and *H. flava*. In 1917, the writer conducted experiments upon *H. fulva*, obtaining results which appear to be of interest if only from a negative standpoint, since the experimental procedure involved seems somewhat similar to the more probable physiological forces at work in the conditions under which the plant forms mature seeds.

Referring to Knuth's Handbook of Flower Pollination, we read that, "according to Sprengel's assertion which Kerner confirms, the plant (*H. fulva*) never sets fruit here, so it is highly probable that in its original home in E. Asia, it is pollinated by such insects as are not to be found in Europe. Maximowicz states that artificial pollination is also ineffective, the flowers do not produce mature seeds in Europe. Sprengel, who pollinated the flowers artificially with their own pollen, also obtained no fruits, etc."

No such limitations affect *H. flava*, indeed Linnaeus believed *H. flava* and *H. fulva* (commonly known as the yellow lily and day lily respectively) to form a composite type species (*H. lilio-asphodelus*), for the genus, and that one was really a variety of the other, a fact readily comprehensible when their great anatomical, if not physiological resemblance, be kept in mind.